




Where can *you* go with KUVAN Powder 500 mg?



KUVAN® (sapropterin dihydrochloride) Powder for Oral Solution 500 mg: a convenient option to help manage your PKU

- For those who respond, KUVAN—in combination with a low-Phe diet—can help keep blood Phe levels low¹
- With KUVAN Powder 500 mg, there are fewer packets to prepare, so you can easily make it part of your daily routine
- Delivers the same Phe-lowering benefits as the 100 mg tablet or packet



Five 100 mg tablets

OR



Five 100 mg packets



One

KUVAN Powder 500 mg packet

Tablets and packets not shown actual size

Taking KUVAN Powder 500 mg is as easy as 1-2-3

KUVAN Powder dissolves completely in 15 seconds in water, apple juice, or a small amount of soft food such as apple sauce or pudding.



1 Open the powder packet



2 Dissolve the powder



3 Drink or eat the mixture within 30 minutes

Get started today. Just ask your clinic team about trying or switching to KUVAN Powder 500 mg.

Experience convenience. Ask for KUVAN Powder 500 mg.

Please see other side for Important Safety Information.

PKU experts recommend

- Treatment should begin at birth and continue throughout life, with blood Phe levels tightly controlled in a range of 2-6 mg/dL or 120-360 $\mu\text{mol/L}$ ²
- Everyone with PKU should be offered a trial of KUVAN*²
- Those who have stopped PKU therapy (including diet and medication) can see improvements in their symptoms if they restart therapy²

BioMarin RareConnections™

There may be 2 co-pays if your prescription includes *both* 500 mg KUVAN Powder *and* 100 mg KUVAN Powder (or 100 mg KUVAN Tablets). For case management and reimbursement support, contact your BPPS representative.

Call: **1.877.MY.KUVAN**
(1.877.695.8826)

Email: support@biomarin-rareconnections.com

*Except for those with two null mutations in *trans*.

Indication

KUVAN® (sapropterin dihydrochloride) Tablets for Oral Use and Powder for Oral Solution are approved to reduce blood Phe levels in people with a certain type of Phenylketonuria (PKU). KUVAN is to be used with a Phe-restricted diet.

Important Safety Information

It is not possible to know if KUVAN will work for you without a trial of the medicine. Your doctor will check your blood Phe levels when you start taking KUVAN to see if the medicine is working.

Starting KUVAN does not eliminate the need for ongoing dietary management. Any change to your diet may impact your blood Phe level. Follow your doctor's instructions carefully. Your doctor and dietitian will continue to monitor your diet and blood Phe levels throughout your treatment with KUVAN **to make sure your blood Phe levels are not too high or too low**. If you have a fever, or if you are sick, your Phe level may go up. Tell your doctor and dietitian as soon as possible so they can make any necessary changes to your treatment.

Children younger than 7 years old treated with KUVAN doses of 20 mg/kg per day are at an increased risk for low levels of blood Phe compared with children 7 years and older. Frequent blood monitoring is recommended in this population to ensure that blood Phe levels do not fall too low.

Tell your doctor if you have ever had liver or kidney problems, have poor nutrition or have a loss of appetite, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

KUVAN is a prescription medicine and should not be taken by people who are allergic to any of its ingredients. KUVAN and other medicines may interact with each other. Tell your doctor about **all the medicines you take**, including prescription and over-the-counter medicines, vitamins, herbal and dietary supplements.

If you forget to take your dose of KUVAN, take it as soon as you remember that day. Do not take 2 doses in a day. If you take too much KUVAN, call your doctor for advice.

The most common side effects reported when using KUVAN are headache, runny nose and nasal congestion, sore throat, diarrhea, vomiting, and cough. Additional adverse reactions reported in connection with worldwide marketing include sore throat, heartburn or pain in the esophagus, inflammation of the lining of the stomach, indigestion, stomach pain, and nausea. These are not all the possible side effects seen with KUVAN. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1.800.FDA.1088.

KUVAN can cause serious side effects, including:

- **Severe allergic reactions.** Stop taking KUVAN and **get medical help right away** if you develop any of these symptoms or a severe allergic reaction:
 - Wheezing or trouble breathing
 - Nausea
 - Flushing
 - Lightheadedness or fainting
 - Coughing
 - Rash
- **Inflammation of the lining of the stomach (gastritis).** Gastritis can happen with KUVAN and may be severe. **Call your doctor right away if you experience:**
 - Severe upper stomach-area discomfort or pain
 - Blood in your vomit or stool
 - Nausea and vomiting
 - Black, tarry stools
- **Too much or constant activity (hyperactivity) can happen with KUVAN.** Tell your doctor if you experience hyperactivity, including fidgeting, moving around or talking too much.

For more information, call BioMarin RareConnections™ at 1.866.906.6100.

Please read the attached Full Patient Information.

B:OMARIN

BioMarin Pharmaceutical Inc.
Copyright 2015. All rights reserved.
US/KUV/0315/0001


KUVAN®
(sapropterin dihydrochloride)
Powder for Oral Solution